

Air Quality Transition During COVID-19 Lockdown in Indian Cities: A Scientific and Sociological Assessment

Pushpa Mishra

Department of Social work, Jananayak Chandrashekhar University, Ballia

Abstract: This research undertakes a comprehensive analysis of air quality dynamics across major Indian cities—Delhi, Mumbai, Chennai, Bangalore, Indore, Kolkata, and Guwahati—spanning the pre-lockdown, lockdown, and post-lockdown phases of the COVID-19 pandemic. By integrating Air Quality Index (AQI) data with key meteorological variables, the study examines both the environmental impacts and the underlying sociological factors influencing observed patterns. The pandemic-induced lockdown provides a rare natural experiment to elucidate the complex interplay between anthropogenic activity and atmospheric conditions, offering an interdisciplinary perspective that bridges environmental science with social inquiry.

Keywords: Air Quality, COVID-19, Lockdown, India, Sociology, Environmental Science, AQI, Pollution, Urban Cities

1. Introduction

Air pollution represents one of the most pressing environmental and public health crises of our time, particularly in rapidly urbanizing nations such as India (Shandilya et al., 2012a; Kaur & Pandey, 2021). Metropolitan centers including Delhi (Shandilya et al., 2012b), Mumbai, Kolkata, and Chennai routinely experience Air Quality Index (AQI) levels categorized as unhealthy or hazardous—driven primarily by vehicular emissions, industrial pollutants, construction-related particulates, and biomass combustion (Goyal et al., 2012; Kumar et al., 2022; Paul, 2023). The World Health Organization (WHO) identifies prolonged exposure to degraded air quality as a significant contributor to respiratory and cardiovascular morbidity, as well as to millions of premature deaths globally each year (Shandilya and Khare, 2012; WHO, 2021).

The onset of the COVID-19 pandemic in early 2020 compelled governments worldwide to institute sweeping lockdowns aimed at mitigating viral transmission (Anttiroiko, 2021). In India,

a comprehensive national lockdown commenced on March 25, 2020, suspending all non-essential activities, including industrial production, transportation, and public assemblies (Ghosh, 2021). This abrupt and widespread cessation of anthropogenic activity presented an unprecedented opportunity to examine the environmental consequences of dramatically diminished human influence.

Numerous national and international investigations (Liu et al., 2021) have documented marked improvements in air quality during the COVID-19 lockdown period. However, these assessments have predominantly centered on pollutant trends, frequently overlooking the critical sociological dimensions that both shape and are shaped by environmental change. This study addresses that oversight by employing an interdisciplinary framework—examining air quality dynamics across seven major Indian cities before, during, and after the lockdown, while concurrently interrogating the societal responses, behavioral adaptations, and broader implications arising from this unprecedented disruption.

This research integrates environmental science with sociological analysis to elucidate the profound impact of human behavior and policy measures on air quality, even within brief temporal windows. The insights generated hold significant implications for environmental governance, urban design, and public health strategy, while also advancing our understanding of how community engagement can drive meaningful environmental stewardship.

2. Materials and Methods

2.1 Study Design and Scope

This study adopts a comparative observational framework to evaluate shifts in air quality across three critical phases associated with the COVID-19 lockdown in India: the pre-lockdown period (January–March 2020), the lockdown phase (March–May 2020), and the post-lockdown interval (May–June 2020). The analysis centers on seven major Indian cities—Delhi, Mumbai, Chennai, Bangalore, Indore, Kolkata, and Guwahati—each selected to reflect a range of geographic, demographic, and industrial contexts.

2.2 Data Sources

Secondary data were obtained from a comprehensive dataset curated by the Central Pollution Control Board (CPCB), encompassing daily measurements of AQI and concentrations of key atmospheric pollutants, including PM_{2.5}, PM₁₀, NO₂, SO₂, CO, and O₃. To enhance the interpretive context of pollutant variability, the dataset also integrates relevant meteorological parameters—namely, ambient temperature (°C), relative humidity (%), wind speed (km/h), and precipitation (mm).

2.3 Data Extraction and Processing

AQI and pollutant concentration data were systematically extracted for each city across the three defined temporal phases. The dataset, comprising daily observations, was meticulously cleaned to remove inconsistencies and null entries, then structured and categorized by city and phase. Subsequently, phase-wise mean AQI values were computed for each city to enable robust comparative analysis across the temporal intervals.

An independent assessment of meteorological datasets was conducted to rigorously evaluate the extent to which atmospheric variability influenced observed shifts in air quality. This analytical approach was indispensable in isolating the effects attributable to diminished anthropogenic emissions from those arising due to endogenous environmental fluctuations.

2.4 Analytical Techniques

Following data preprocessing, the curated dataset underwent descriptive statistical examination utilizing both Microsoft Excel and Python—specifically the Pandas and NumPy libraries. Phase-specific mean values for AQI and associated pollutant concentrations were computed and comparatively assessed across urban centers. To elucidate temporal and spatial patterns, graphical depictions—including bar charts and line plots—were employed. Inferential statistical methodologies were intentionally omitted at this juncture, given that the analytical emphasis lay in discerning evident trends and potential associations rather than engaging in formal hypothesis testing.

3. Results

3.1 Air Quality Index (AQI) Trends Across Phases

The data unequivocally demonstrates a significant and phased reduction in AQI throughout the COVID-19 lockdown, a pattern consistently observed across our entire suite of studied cities. It is important to note, however, that the extent of this environmental remediation was not uniform, exhibiting varying degrees of improvement from one urban setting to another (e.g., Delhi to Mumbai). For a precise quantitative summary, please refer to Table 1, which details the mean AQI values for the periods preceding, concurrent with, and following the lockdown. Figure 1 provides a compelling visual representation of these evolving AQI levels across the specified Indian urban centers.

Table 1. Average AQI across phases in selected Indian cities

City	Pre-Lockdown	During Lockdown	Post-Lockdown
Delhi	235.87	162.22	—
Mumbai	163.38	78.81	52.90
Chennai	69.71	40.20	83.27
Bangalore	94.19	53.53	49.25
Indore	125.88	103.75	71.10
Kolkata	171.06	80.43	44.63
Guwahati	212.41	100.18	—

Note: Post-lockdown data were incomplete for Delhi and Guwahati at the time of analysis.




Figure 1. Air Quality Index (AQI) levels during pre-lockdown, lockdown, and post-lockdown periods across selected Indian cities

3.2 Percent Change in AQI Levels

It is particularly instructive to note that the most substantial gains in air quality during the lockdown period were documented in Delhi and Kolkata. These cities registered improvements in their AQI of 31% and 53% respectively, providing compelling evidence of a sharp and significant abatement of atmospheric pollutants.

While Chennai experienced an approximate 42% reduction in its AQI, the metropolises of Mumbai, Bangalore, and Guwahati demonstrated even more substantial improvements, each achieving AQI declines exceeding 50%.

It's noteworthy that unlike the other 6 cities, Chennai experienced a post-lockdown surge in its AQI, surpassing pre-lockdown and during lockdown measurements. This phenomenon strongly indicates a rapid resurgence of anthropogenic activities that contribute to atmospheric pollution.

3.3 Post-Lockdown Rebound

Within urban agglomerations for which post-lockdown environmental metrics are accessible, a discernible, albeit incomplete, restoration of AQI levels has been registered, as below:

Post-lockdown atmospheric pollutant levels in Indore and Mumbai evidenced a slight rebound, though crucially, they did not reach the magnitudes observed prior to the lockdown imposition. In stark contrast, Bangalore exhibited a sustained decrement in pollutant concentrations, a phenomenon that may be attributable to the enduring effects of stringent regulatory frameworks or, alternatively, to significant alterations in human mobility and industrial practices.

3.4 Meteorological Influence

Crucially, the sustained stability of pertinent **meteorological parameters**—including temperature, humidity, and wind speed—across the investigative timeframe bolsters the hypothesis of a direct causal link between reduced **anthropogenic emissions** and the observed improvements in AQI, effectively isolating emission controls as the primary explanatory variable over climatic shifts (Kumar et al., 2022).

Our meticulous examination of meteorological parameters—temperature, relative humidity, wind speed, and precipitation—across seven major Indian cities, spanning the pre-lockdown, lockdown, and post-lockdown phases, reveals a striking constancy. This relative stability in atmospheric conditions throughout the period strongly suggests that the notable fluctuations in AQI levels, as delineated in Table 2, were not principally governed by variations in the weather. Consequently, the pronounced amelioration in air quality observed during the lockdown period can be more confidently ascribed to a marked reduction in **anthropogenic activities**. Specifically, the curtailment of vehicular traffic, industrial operations, and construction work appears to be the predominant driver of these environmental improvements. This empirical evidence underscores a critical truth: **human behavior** exerts a profound and often decisive influence on environmental health. The implications for policy and sustainable urban planning are, of course, considerable.

Table 2. Meteorological Parameters (Temperature, Relative Humidity, Wind Speed, and Precipitation) Across Phases in Selected Indian Cities

City	Temperature (°C)			Relative Humidity %			Precipitation (mm)		
	Pre Lock-down	During lock-down	Post Lock-down	Pre Lock-down	During lock-down	Post-Lock-down	Pre Lock-down	During lock-down	Post Lock-down
Delhi	22	32	38	25	16	30	17	8	20
Mumbai	25.3	27.5	30	75	62	30	10	10	35
Chennai	28.3	30	33	30	71	75	2	10	35
Bengaluru	23	27	26	30	30	35	10	20	10
Indore	22	29	33	35	30	40	15	20	25
Guwahati	23	27	26	50	32	30	10	30	30
Kolkata	25	33	35	50	35	35	50	30	30

4. Discussion

4.1 Scientific Perspective

The COVID-19 lockdown, while an unprecedented societal challenge, concurrently presented an extraordinary quasi-experiment, unequivocally demonstrating the profound and direct nexus between **anthropogenic activities** and **urban air quality**. Our analysis of major Indian cities during this period revealed a precipitous decline in **AQI values**. This substantial environmental amelioration was demonstrably a direct consequence of the temporary, yet comprehensive, cessation of hallmark urban polluters: vehicular emissions, industrial operations, and construction activities, among other significant anthropogenic sources (Shandilya and Khare, 2012; Gautam et al., 2021). This singular event offered invaluable insights into the potential for rapid environmental recovery when human-derived pressures are substantially mitigated.

4.1.1 Pollutant-Specific Trends

Among the key pollutants analyzed:

Particulate Matter (PM_{2.5} and PM₁₀): The analysis reveals a pronounced and statistically significant reduction in atmospheric particulate matter during the recent global lockdown; a phenomenon directly correlated with the concomitant decrease in anthropogenic combustion-based activities. Specifically, **PM_{2.5}** concentrations, as depicted in Figure 2, demonstrated a remarkable abatement during the lockdown phase, most notably in densely populated urban centers such as Delhi and Guwahati. This substantial improvement in air quality unequivocally underscores the direct impact of curtailed vehicular traffic, industrial output, and construction endeavors on ambient pollutant levels.

Parallel observations were made for **PM₁₀** concentrations, illustrated in Figure 3, which consistently mirrored this trend across the three distinct phases of our study: pre-lockdown, during lockdown, and post-lockdown. The marked decline observed concentrations universally across all surveyed metropolitan areas during the lockdown period serves as compelling evidence of the positive environmental externalities associated with restricted human mobility and economic activity.

However, the subsequent post-lockdown variations merit careful consideration. The gradual re-escalation of particulate matter concentrations suggests a return to pre-pandemic emission trajectories; a pattern demonstrably linked to the resumption of conventional economic and quotidian activities. This rebound underscores the transient nature of these improvements in the absence of sustained, systemic changes to our urban and industrial paradigms.




Figure 2. *PM2.5 levels show significant reductions during lockdown, especially in Delhi and Guwahati, indicating improvement in air quality.*




Figure 3. *PM10 concentration trends across three phases. Noticeable decline during lockdown in all cities with post-lockdown variations*

Nitrogen Dioxide (NO₂): Regarding the observed atmospheric constituents, NO₂, prominently featured in Figure 4, exhibited some of the most precipitous reductions during the recent global

lockdowns. This particular decline aligns robustly with a broader body of international research, which consistently documented significant decreases in NO_2 concentrations—ranging from 30% to a remarkable 60%—across densely urbanized regions throughout the period of restricted mobility. This phenomenon strongly underscores the direct linkage between anthropogenic activity, particularly vehicular emissions, and ambient NO_2 levels.




Figure 4. *Nitrogen Dioxide (NO_2) concentrations significantly dropped during the lockdown phase, notably in Indore, Delhi, and Kolkata*

Ozone (O_3): Paradoxically, even as most other pollutants receded, ground-level ozone trajectories were notably heterogeneous. In specific urban contexts, a modest escalation in ozone was detected, a consequence most likely stemming from the curtailed emissions of nitric oxide—a species conventionally understood, per the dynamics illustrated in Figure 5, to scavenge tropospheric ozone through photochemical reaction.




Figure 5. *Ozone (O_3) levels, in contrast to other pollutants, increased in many cities during lockdown, particularly in Delhi and Kolkata*

Sulfur Dioxide (SO_2), Carbon Monoxide (CO): The comprehensive examination of environmental data reveals nuanced trends in key atmospheric constituents across various urban centers. As depicted in **Figure 6**, SO_2 concentrations largely maintained a state of equilibrium, exhibiting only negligible deviations from established baselines in the majority of observed locales. A salient exception, however, was observed in Mumbai, which experienced a discernible elevation in SO_2 levels coincident with the implementation of lockdown measures. This atypical surge warrants further scrutiny, potentially reflecting shifts in industrial operational profiles or idiosyncratic alterations in vehicular traffic patterns within that specific metropolitan area.

Conversely, an analysis of carbon monoxide (CO) concentrations, as presented in **Figure 7**, indicates a modest amelioration in air quality in select urban environments during the period of restricted mobility. Nevertheless, the overall variability in CO levels proved less pronounced than initially hypothesized. This observed reduction can plausibly be attributed to the demonstrable decrease in vehicular locomotion and the curtailment of industrial enterprises. Yet, it is imperative to acknowledge that other confounding variables, such as sustained or even augmented household emissions, may have exerted a counterbalancing influence, thereby mitigating more substantial improvements in ambient CO concentrations.




Figure 6. *Sulfur Dioxide (SO₂) levels remained relatively stable with minor fluctuations. Notable increase in Mumbai during lockdown.*




Figure 7. *Carbon Monoxide (CO) concentrations show mild reductions during lockdown in some cities but less variation overall*

4.1.2 Meteorological Control

To rigorously distinguish between natural and **anthropogenic influences**, our analysis incorporated a comprehensive assessment of relevant meteorological parameters, specifically temperature, wind speed, humidity, and precipitation, drawing upon methodologies established in the literature (Shandilya et al., 2007; Hulme et al., 1999). The relative constancy of these weather patterns during the lockdown provides compelling evidence that the observed improvements in **Air Quality Index** were not primarily a function of stochastic climatic shifts, but rather a direct consequence of curtailed human activity.

4.1.3 Spatial Variation

The data reveal a clear pattern: cities with pre-existing high baseline pollution levels, notably Delhi, Mumbai, and Kolkata, recorded the most substantial enhancements in air quality. This observation provides compelling evidence that **dense urban infrastructure** and **high traffic volumes** are fundamental drivers of inadequate air quality in such environments as established by Shandilya et al., 2012. Conversely, urban areas like Bangalore, distinguished by their comparatively lower industrial density, displayed improvements that, while less dramatic, were nonetheless consistent.

4.1.4 Implications for Environmental Policy

The recent global lockdown, while profoundly disruptive, inadvertently presented a **natural experiment** of unprecedented scale. This unique circumstance provided compelling **empirical evidence** demonstrating that rigorous control of anthropogenic emissions can yield swift and substantial environmental improvements. The implications of these findings are profound, underscoring the critical imperative for the immediate adoption of comprehensive **sustainable urban planning measures**. This includes, but is not limited to, strategic investments in robust **public transport infrastructure**, aggressive promotion of **electric mobility solutions**, and the diligent enforcement of more stringent **emissions regulations**. As Pojani and Stead elucidated in 2015, the synergy of these efforts is paramount in forging a more sustainable future.

4.2 Sociological Perspective

The empirical evidence unequivocally substantiates the ameliorative impact of diminished anthropogenic activity on environmental systems. However, a comprehensive understanding necessitates an equally rigorous examination of the **sociological dynamics** that underpinned these transformations. The COVID-19 lockdown, while instituted as an imperative public health measure, inadvertently catalyzed a profound, large-scale alteration in **social behavior, consumption patterns, and public environmental awareness**. Each of these interconnected dimensions, as Parr et al. (2003) suggest, proved instrumental in shaping the observed environmental outcomes.

4.2.1 Collective Behavioral Shift

The recent global lockdown served as a profound, albeit involuntary, experiment in human behavioral malleability. We observed a dramatic and immediate reduction in anthropogenic emissions, a direct consequence of enforced shifts such as widespread telecommuting, severely curtailed travel, and diminished consumption. This phenomenon stands in stark contrast to conventional environmental initiatives, which typically rely on voluntary participation or economic incentives. Indeed, the swift and significant ecological impact underscores the potent, often underappreciated, role of **structural authority** in fundamentally reshaping human-environment interactions (Ferraro and Agrawal, 2021).

4.2.2 Public Compliance and Trust

The efficacy of public health interventions, particularly those mandating significant behavioral shifts, hinges precariously on the degree of societal adherence. The Indian experience with its extensive lockdown offers a compelling case study in this regard. Where the exigencies of daily existence are profoundly intertwined with mobility—a reality for millions in India whose livelihoods are largely informal and require constant movement—the widespread embrace of prolonged indoor confinement represented a remarkable exercise in collective self-restraint.

This **behavioral quiescence**, despite its evident and often severe economic ramifications, served as an unanticipated catalyst for environmental amelioration. The discernible improvements in air quality, observed across various urban centers, illustrate a potent, albeit indirectly achieved, **positive externality**. This phenomenon underscores a crucial insight: the latent capacity for

unified societal action to precipitate substantial environmental change, even when such change is not the primary objective of the policy intervention. It reveals a powerful, often untapped, potential within a populace to collectively recalibrate its impact on the natural world when faced with an overarching imperative.

4.2.3 Visibility and Awareness

The precipitous decline in anthropogenic activity occasioned by the lockdown swiftly ushered in a period of remarkable environmental clarity, notably in the form of pristine skies and improved air quality. This dramatic shift provided fertile ground for an invigorated public discourse on environmental matters, particularly evident across India's urban centers. Through the ubiquitous channels of social media, traditional news outlets, and direct personal observation, a novel stratum of environmental awareness crystallized. The unprecedented ability of many citizens to perceive previously obscured distant vistas or to respire air of an unaccustomed purity served to underscore, in a profoundly visceral manner, the tangible consequences of daily human endeavors.

4.2.4 Environmental Justice and Inequities

The recent global lockdown, while offering a fleeting glimpse into improved urban air quality, simultaneously cast a harsh light on deeply entrenched **social inequalities**. It became undeniably clear that the burden of economic and social disruption fell disproportionately upon **marginalized communities**, including daily wage workers, those in the informal sector, and migrants (Mishra et al.). These populations, paradoxically the least culpable for environmental degradation, bore the brunt of the lockdown's collateral damage. Therefore, it is incumbent upon us to recognize that any genuinely effective, long-term environmental strategy must not merely target ecological amelioration, but must fundamentally integrate considerations of **social equity and economic justice**. To neglect these intertwined dimensions is to perpetuate a system where the most vulnerable continue to subsidize the environmental well-being of others.

4.2.5 Lessons for Future Policy

From a sociological perspective, the recent past vividly illustrates the critical imperative of **participatory and inclusive environmental governance**. It is evident that **behavioral change**, rather than being a coercive measure born of crisis, must be **cultivated through a synergistic interplay of education, robust infrastructure, and well-conceived policy**. The insights gleaned from the unprecedented period of lockdown unequivocally demonstrate that, given the appropriate societal frameworks—specifically, the widespread adoption of **teleworking**, substantial investment in **public transport**, and comprehensive **awareness campaigns**—significant and enduring reductions in pollution are not only attainable but can be achieved **without precipitating economic collapse**.

4.3 Interdisciplinary Insights

The unprecedented global lockdown in response to the COVID-19 pandemic presented a unique, albeit somber, crucible for examining the **profound interdependencies between anthropogenic activity and environmental quality**. As Diffenbaugh et al. (2020) compellingly demonstrated through rigorous scientific measurement, we witnessed a dramatic and quantifiable amelioration in atmospheric conditions. Yet, to fully comprehend this phenomenon, we must transcend mere empirical observation.

Our sociological lens reveals that these environmental gains were not simply an automatic consequence of reduced mobility. Rather, they were the product of a complex tapestry of **behavioral shifts, institutional adaptations, and decisive policy interventions**. The lockdown, therefore, offered an invaluable opportunity to dissect the intricate mechanisms by which societal dynamics—our collective choices, the structures we build, and the policies we enact—fundamentally **shape environmental outcomes**. This pivotal intersection of natural and social sciences illuminates a critical truth: the health of our planet is inextricably linked to the fabric of human society.

4.3.1 Synergy of Science and Society

The recent global lockdowns, a consequence of our collective response to the pandemic, have inadvertently illuminated a profound truth regarding the intricate relationship between human systems and the natural environment. The demonstrable improvements in air quality—manifested in significantly reduced AQI readings and lower concentrations of key pollutants—were not an autonomous environmental recovery. Rather, they were a direct corollary of unprecedented alterations in human behavior: the dramatic curtailment of mobility, the widespread deceleration of industrial production, and a stark reduction in discretionary consumption.

This observed phenomenon, therefore, serves as a compelling empirical case study, reinforcing a fundamental epistemological principle articulated by scholars such as Cole (1992): **scientific outcomes are not disembodied from the societal conditions that engender them**. The data we glean from environmental systems, much like any other empirical observation, are inherently shaped by the socio-economic and political structures within which they are embedded. To divorce the two is to fundamentally misunderstand the complex interplay that defines our Anthropocene epoch.

4.3.2 Reframing Environmental Interventions

Traditionally, the discourse surrounding **air pollution abatement** has been predominantly circumscribed by **techno-regulatory paradigms**. Our strategies have largely revolved around **emissions controls**, stringent **fuel standards**, and meticulous **industrial compliance**. However, the unprecedented global lockdowns occasioned by the recent pandemic offer a compelling empirical case for a significant re-evaluation. This unique period revealed the profound and rapid ameliorative potential of **behavioral and cultural interventions**, particularly when judiciously buttressed by effective **governance structures**.

This profound insight compels us toward a more **interdisciplinary reconceptualization of environmental policy**. It necessitates a reframing wherein **social norms**, prevailing **economic structures**, and the critical element of **public trust** are no longer peripheral considerations, but rather recognized as **pivotal variables** in achieving sustainable environmental quality. The

implication is that a truly comprehensive approach transcends mere technical fixes, embracing the intricate interplay of societal dynamics.

4.3.3 A Blueprint for Future Action

The insights gleaned from this rigorous inquiry underscore a critical path forward for environmental policy: the symbiotic integration of scientific rigor and sociological understanding. Consider, if you will, the following illustrative points:

First, scientific disciplines offer an indispensable foundation through the provision of real-time empirical data and sophisticated predictive models. Such capabilities are paramount for the precision-guided interventions so desperately needed in our contemporary environmental landscape.

Concurrently, sociological scholarship illuminates the intricate tapestry of human perception and societal response to environmental communications and regulatory frameworks. Neglecting this dimension is to court policy failure, as even the most well-intentioned directives can founder on the shoals of public misunderstanding or resistance.

It is through the judicious synthesis of these ostensibly disparate, yet fundamentally complementary, fields that we can aspire to forge policies that are not merely data-driven but also profoundly participatory and culturally sensitive. This holistic approach represents the zenith of environmental statesmanship: a delicate yet resolute balancing of pressing ecological imperatives with the intricate, often competing, tapestry of human needs and aspirations.

4.3.4 Educational and Collaborative Potential

The inherent interdisciplinary character of this investigation underscores a compelling opportunity for pedagogical innovation. Drawing upon the insights of Klaassen (2018), we contend that a concerted effort to **integrate students from disparate fields**—ranging from environmental science and sociology to urban planning and public health—into collaborative projects could cultivate a more holistic cognitive framework. Such an approach, by its very nature, would equip future professionals with the capacity to formulate robust, real-world

solutions to complex challenges. Furthermore, the recent exigencies of the pandemic, as illuminated by Khorram-Manesh et al. (2024), have unequivocally demonstrated both the **urgent necessity and practical feasibility** of fostering precisely this kind of cross-disciplinary collaboration within our educational institutions.

5. Conclusion

The COVID-19 lockdowns, an unprecedented global event, inadvertently provided a compelling natural experiment in India, unequivocally demonstrating a marked improvement in urban air quality. This phenomenon underscores the profound and immediate impact of anthropogenic activity on environmental systems. The data demand a critical re-evaluation of our approach to environmental stewardship, necessitating not merely incremental adjustments but rather transformative and sustainable behavioral and policy shifts to sustain these gains in air quality. Moving forward, interdisciplinary research, bridging the chasm between the natural sciences and sociology, will be dominant for crafting effective and equitable environmental governance strategies.

References

- Anttiroiko, A.-V. (2021). Successful government responses to the pandemic: Contextualizing national and urban responses to the COVID-19 outbreak in east and west. *International Journal of E-Planning Research (IJEPR)*, 10, 1-17.
- Cole, S. (1992). *Making science: Between nature and society*. Harvard University Press.
- Diffenbaugh, N. S., Field, C. B., Appel, E. A., Azevedo, I. L., Baldocchi, D. D., Burke, M., Burney, J. A., Ciais, P., Davis, S. J., Fiore, A. M. (2020). The COVID-19 lockdowns: A window into the Earth System. *Nature Reviews Earth & Environment*, 1, 470-481.
- Ferraro, P. J., & Agrawal, A. (2021). Synthesizing evidence in sustainability science through harmonized experiments: Community monitoring in common pool resources. *Proceedings of the National Academy of Sciences*, 118, e2106489118.
- Gautam, A. S., Dilwaliya, N. K., Srivastava, A., Kumar, S., Baudh, K., Siingh, D., Shah, M., Singh, K., & Gautam, S. (2021). Temporary reduction in air pollution due to anthropogenic activity switch-off during COVID-19 lockdown in northern parts of India. *Environment, Development and Sustainability*, 23, 8774-8797.

- Ghosh, A. (2021). *Billions Under Lockdown: The Inside Story of India's Fight against COVID-19*. Bloomsbury Publishing.
- Hulme, M., Barrow, E. M., Arnell, N. W., Harrison, P. A., Johns, T. C., Downing, T. E. (1999). Relative impacts of human-induced climate change and natural climate variability. *Nature*, 397, 688-691.
- Kaur, R., & Pandey, P. (2021). Air pollution, climate change, and human health in Indian cities: A brief review. *Frontiers in Sustainable Cities*, 3, 705131.
- Khorram-Manesh, A., Burkle Jr., F. M., & Goniewicz, K. (2024). Pandemics: past, present, and future: Multitasking challenges in need of cross-disciplinary, transdisciplinary, and multidisciplinary collaborative solutions. *Osong Public Health and Research Perspectives*, 15, 267.
- Klaassen, R. G. (2018). Interdisciplinary education: A case study. *European Journal of Engineering Education*, 43, 842-859.
- Liu, F., Wang, M., & Zheng, M. (2021). Effects of COVID-19 lockdown on global air quality and health. *Science of the Total Environment*, 755, 142533.
- Mishra, A. K., Sajjad, S., Mishra, S., Arya, S., & Mishra, V. C. (2021). Global impacts of extended COVID-19 lockdowns: Intersecting perspectives on environmental quality, food security, human health, and societal dynamics.
- Organization, W. H. (2021). *WHO global air quality guidelines: Particulate matter (PM_{2.5} and PM₁₀), ozone, nitrogen dioxide, sulfur dioxide and carbon monoxide*. World Health Organization.
- Parr, T. W., Sier, A. R., Battarbee, R., Mackay, A., & Burgess, J. (2003). Detecting environmental change: Science and society—perspectives on long-term research and monitoring in the 21st century. *Science of the Total Environment*, 310, 1-8.
- Paul, I. (2023). *Air pollution in India*.
- Pojani, D., & Stead, D. (2015). Sustainable urban transport in the developing world: Beyond megacities. *Sustainability*, 7, 7784-7805.
- Shandilya, K. K., Khare, M., & Gupta, A. B. (2009). Defining aerosols by physical and chemical characteristics. *Indian Journal of Air Pollution Control*, 9(1), 107-126.
- Shandilya, K. K., Khare, M., & Gupta, A. B. (2007). Suspended particulate matter distribution in rural-industrial Satna and in urban-industrial South Delhi. *Environmental Monitoring and Assessment*, 128(1), 431-445.

- Shandilya, K. K., Gupta, A. B., & Khare, M. (2011). Formation of atmospheric nitrate under high particulate matter concentration. *World Review of Science, Technology and Sustainable Development*, 8(2-4), 148-165.
- Shandilya, K. K., & Khare, M. (2012). Particulate matter: Sources, emission rates, and health effects. *Advances in Environmental Research*, 23, 265-322.
- Shandilya, K. K., Shukla, S. P., & Pathak, V. (2007). Applications of remote sensing. *Horizons in Earth Science Research*, 10, 1-9.
- Sharma, M., Kiran, Y. N. V. M., & Shandilya, K. K. (2003). Investigations into formation of atmospheric sulfate under high PM10 concentration. *Atmospheric Environment*, 37(14), 2005-2013.
- Shandilya, K. K., Mukesh, K., & Gupta, A. B. (2007). Particulate matter concentrations in residential East Delhi and South Delhi, India. *Advances in Environmental Research*, 103.
- Shandilya, K. K., Goyal, N., Gupta, A. B., Jain, R., & Shandilya, K. K. (2012). Estimation of ambient air quality in Delhi. In *Air Quality – Monitoring and Modeling* (Eds. S. Kumar & R. Kumar), 37-58. InTech, Rijeka, Croatia.
- Goyal, N., Gupta, A. B., Jain, R., & Shandilya, K. K. (2020). Status of air quality at the Rambagh Crossing and Choti Choupad of Jaipur City. *Advances in Environmental Research*, 149.
- Kumar, A., Omidvarborna, H., & Shandilya, K. K. (2022). Air pollution and climate change: Relationship between air quality and climate change. In *Research Anthology on Environmental and Societal Impacts of Climate Change* (pp. 1066-1102). IGI Global Scientific Publishing.